

The background of the slide features a blurred image of laboratory glassware, including vials and bottles, overlaid with a large blue diagonal shape.

NOVEL THERAPEUTICS  
FOR **METABOLIC &**  
**ENDOCRINE DISORDERS**

Corporate Presentation

May 2019

# Forward-Looking Statements

This presentation contains statements about our future expectations, plans and prospects that constitute forward-looking statements for purposes of the safe harbor provisions of the Private Securities Litigation Reform Act of 1995. Actual results may differ materially from those indicated by these forward-looking statements as a result of various important factors, including risks relating to: both our and our collaborators' ability to successfully research, obtain regulatory approvals for, develop and commercialize products based upon our technologies; our ability to obtain and maintain proprietary protection for our technologies and product candidates; our reliance on third parties to manufacture our preclinical and clinical drug supplies; competitive pressures; our ability to obtain and maintain strategic collaborations; compliance with our in-license agreements; our ability to successfully execute on, and receive favorable results from, our proprietary drug development efforts; market acceptance of our drug candidates; retaining members of our senior management; and our ability to raise additional funds to finance our operations.

The forward-looking statements included in this presentation represent our views as of the date of this presentation. We anticipate that subsequent events and developments will cause our views to change. While we may elect to update these forward-looking statements in the future, we specifically disclaim any obligation to do so. These forward-looking statements should not be relied upon as representing our views as of any date subsequent to the date of this presentation.

For more information regarding risks and uncertainties that could affect the results of our operations or financial condition review our filings with the Securities and Exchange Commission (in particular, our most recent Annual Report on Form 10-K and any subsequently filed Quarterly Reports on Form 10-Q).

# Investment Highlights

## ► Focused on best-in-class drugs for metabolic and endocrine diseases

- Two Phase 2 programs supported by positive clinical data

## ► Metabolic Disease Program: VK2809 for NASH

- Novel, selective thyroid receptor- $\beta$  (TR $\beta$ ) agonist
- Phase 2 results demonstrate significant reduction in liver fat content, lipids

## ► Rare Disease Program: VK0214 for X-ALD

- Second selective thyroid receptor- $\beta$  agonist
- In vivo data show improvement in key biomarkers

## ► Musculoskeletal Program: VK5211 for hip fracture recovery

- Non-steroidal selective androgen receptor modulator (SARM)
- Phase 2 results demonstrate significant increases in lean body mass

# Pipeline Overview

| Product Candidates             | Indication                   | Development Stage |         |         |         | Status                                                 |
|--------------------------------|------------------------------|-------------------|---------|---------|---------|--------------------------------------------------------|
| Development Programs           |                              | Preclin           | Phase 1 | Phase 2 | Phase 3 |                                                        |
| VK2809<br>(TR $\beta$ agonist) | NASH                         |                   |         |         |         | Initiate clinical study in biopsy-confirmed NASH, 2H19 |
| VK0214<br>(TR $\beta$ agonist) | X-ALD                        |                   |         |         |         | IND planned, 2019                                      |
| Other Programs                 |                              | Preclin           | Phase 1 | Phase 2 | Phase 3 |                                                        |
| VK5211<br>(SARM)               | Hip fracture, muscle wasting |                   |         |         |         | Phase 2 completed                                      |
| VK0612<br>(FBPase inhibitor)   | Type 2 Diabetes              |                   |         |         |         | Phase 2a completed                                     |
| VK1430<br>(DGAT-1 inhibitor)   | Hypertriglyceridemia, NASH   |                   |         |         |         | Preclinical                                            |



# Metabolic Disease Program


VK2809: Selective Thyroid Receptor- $\beta$  Agonist

Liver Disorders

# Metabolic Disease Program: Selective Thyroid- $\beta$ Agonists

- ▶ **Proprietary platform for small molecule thyroid hormone mimetics**
  - Highly tissue and receptor selective
  - Produce potent lipid reductions in animals and humans
  - Unique chemical scaffolds, expected wider safety window vs. other approaches
- ▶ **Biological profiles suggest potential benefit in multiple indications**
  - Broad: NASH, hypercholesterolemia, dyslipidemia
  - Rare: X-linked adrenoleukodystrophy (X-ALD), other
- ▶ **Lead molecules VK2809, VK0214**
  - Oral, once-daily formulations
  - VK2809: Successfully completed Phase 2 trial in NAFLD and hypercholesterolemia
  - VK0214: Preclinical, X-ALD

# Thyroid Receptor Overview




Key steps in receptor activation: Endogenous thyroid hormone T<sub>3</sub> crosses mitochondrial membrane (1) binding thyroid receptor TR, dissociating co-repressor CoR (2). Subsequent binding of co-activator CoA (3) results in altered gene expression (4). RXR: retinoid X receptor; TRE: thyroid response element.

- ▶ Nuclear hormone receptors
- ▶ Two major subtypes
  - **Thyroid beta receptor:**  
Liver, brain; modulates cholesterol, triglyceride levels
  - **Thyroid alpha receptor:**  
Cardiac tissue, modulates heart rate, contraction

Therapeutic goal, lipid setting:  $\beta$ -receptor selectivity;  
minimize alpha-effects


# VK2809: Unique Liver-Targeted Characteristics




VK2809, Novel Prodrug

Following oral dosing:

- Cyp3A4-mediated cleavage of prodrug
- 3A4 primarily expressed in liver
- Results in targeted delivery of drug to liver




VK2809A, Potent TR $\beta$  Agonist,  
2.2 nM Ki



VK2809A

- ▶ 17:1 selective for  $\beta$ : $\alpha$
- ▶ Highly negatively charged
  - Poor passive diffusion
- ▶ Not actively transported
  - Due to altered chemistry
- ▶ Targeted hepatic re-uptake
  - Selective liver re-absorption via hepatic OATP1B3 transporter




T3 Thyroid Hormone

- ▶ 1:2 selective for  $\beta$ : $\alpha$
- ▶ Effectively neutral charge
- ▶ Active uptake in multiple tissues via MCT8
- ▶ Broad systemic availability
- ▶ Impractical for development due to safety

Selective activation, differentiated chemistry lends VK2809 liver selectivity; potentially minimizes risk of systemic effects




# VK2809: Liver-Selective Transcriptional Effects



VK2809 shows minimal effects on gene expression in extrahepatic tissues

# Thyroid Receptor $\beta$ Agonists for NAFLD and NASH

## Development of NASH:




- ▶  $\beta$ -Receptor: Key role in lipid metabolism; systemic and liver-specific effects
- ▶ Receptor localized to liver, limited ex-hepatic expression
- ▶ In vivo evidence suggests  $\beta$ -activation provides anti-fibrotic benefits
- ▶ Clinical data indicate a correlation between reduced liver fat, improvement in NAS

An agent that reduces liver fat, improves systemic lipids, and antagonizes fibrotic signaling could provide multi-pronged benefits in NASH

# VK2809 Significantly Reduces Steatosis in Diet-Induced NASH

- ▶ Evaluation in biopsy-confirmed diet-induced NASH model
  - Rodent model designed to reflect progression of disease in humans
  - Animals biopsied pre-study; only those with NASH *and* fibrosis selected
  - VK2809 dosed once-daily for 8 weeks

## Change in Liver Lipids Following 8 Weeks Dosing With VK2809




|               |         |         |         |         |
|---------------|---------|---------|---------|---------|
| % Difference: | -70.0%  | -64.6%  | -79.5%  | -39.7%  |
| p-value:      | <0.0001 | <0.0001 | <0.0001 | <0.0001 |

Treatment with VK2809 significantly improves lipids, steatosis, NAS at 8 weeks; well-tolerated with no evidence of toxicity

# VK2809 Improves Fibrosis in Diet-Induced NASH Model

- ▶ Significant reductions in fibrosis, collagen, hydroxyproline after 8 weeks
- ▶ Supports thesis that selective TR $\beta$  activation produces broad metabolic benefits
- ▶ Gene expression data show improvements in genes related to lipid metabolism, insulin sensitivity, fibrogenic signaling

## Change in Liver Fibrosis Following 8 Weeks Dosing With VK2809




|               |        |        |        |
|---------------|--------|--------|--------|
| % Difference: | -50.2% | -60.2% | -46.3% |
| p-value:      | <0.01  | <0.005 | 0.01   |

VK2809 significantly improved NASH and fibrosis in this model

# VK2809: Representative Gene Effects, DIO NASH Model

- ▶ VK2809 reduces expression and signaling of key fibrosis drivers
- ▶ Improvement in genes associated with lipid metabolism, insulin sensitivity also observed
- ▶ Overall gene expression changes align with observed improved histology

## Change Pro-Fibrogenic Gene Expression Following VK2809 Treatment




|              |        |        |        |        |        |
|--------------|--------|--------|--------|--------|--------|
| % Difference | -27.1% | -36.3% | -37.0% | -56.3% | -64.7% |
| p-value      | 0.07   | <0.05  | <0.05  | <0.001 | <0.001 |

TR $\beta$  mechanism provides broad histologic benefits; effects span steatosis, inflammation, fibrosis.

# VK2809 Clinical Highlights: 14-Day Phase 1b Study

- ▶ Placebo-controlled trial (n=56), mild hypercholesterolemia
  - Enrollment criterion: LDL  $\geq$  100 mg/dL
- ▶ Results: clinically, statistically significant reductions in LDL
- ▶ Up to 79% reduction in triglycerides
  - Major component of liver fat
- ▶ Encouraging safety and tolerability, no SAEs




**Placebo-adjusted reduction, LDL:**

|         |          |          |          |
|---------|----------|----------|----------|
| -15.2%  | -27.1%   | -41.2%   | -36.6%   |
| p=0.026 | p=0.0003 | p<0.0001 | p<0.0001 |

**Placebo-adjusted reduction, triglycerides:**

|         |          |          |          |
|---------|----------|----------|----------|
| -34.8%  | -61.0%   | -62.1%   | -78.6%   |
| p=0.052 | p=0.0019 | p=0.0007 | p=0.0001 |

# VK2809-201: Phase 2 Study Design




- ▶ Multi-arm, dose-ranging, 12 week Phase 2 trial
  - Target enrollment: 20 patients per arm
  - **Primary endpoint:** Change in LDL-C vs. placebo
  - **Secondary endpoint:** Change in liver fat by MRI-PDFF
  - **Exploratory endpoints:** Changes in atherogenic proteins

# VK2809 Significantly Reduced LDL-C After 12 Weeks

- ▶ All VK2809 cohorts statistically significantly reduced vs. baseline
- ▶ Placebo-adjusted change from baseline
  - 5 mg QD: -23.7 mg/dL
  - 10 mg QOD: -27.1 mg/dL
  - 10 mg QD: -28.3 mg/dL


## Mean % Change in LDL-C at 12 Weeks



# VK2809 Produced Significant Relative Reductions in Liver Fat

- ▶ Significant Relative Reductions from Baseline in Liver Fat by MRI-PDFF
- ▶ Maximal reductions at Week 12
  - 5 mg QD: 78%
  - 10 mg QOD: 72%
  - 10 mg QD: 76%

## Median Relative % Change in Liver Fat at 12 Weeks




|                     |       |        |        |        |
|---------------------|-------|--------|--------|--------|
| % Change            | -9.4% | -53.8% | -56.5% | -59.7% |
| p-value vs. placebo | -     | 0.0001 | 0.0018 | 0.0004 |

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001

# VK2809 Produced Significant Absolute Reductions in Liver Fat

- ▶ Significant Absolute Reductions from Baseline in Liver Fat by MRI-PDFF
- ▶ Maximal reductions at Week 12
  - Placebo: -6%
  - 5 mg QD: -18%
  - 10 mg QOD: -21%
  - 10 mg QD: -19%

## Mean Absolute % Change in Liver Fat at 12 Weeks



% Change (SD)

-1.1%  
(2.8)

-8.7%  
(4.8)

-8.9%  
(6.2)

-10.6%  
(5.2)

p-value vs. placebo

-

0.0014

0.013

0.0030

# Representative Fat Reduction, VK2809 and Placebo Subject




## Overall Mean Hepatic Fat Values

| Subject, Dose      | Baseline | Week 12 | Absolute Change | Relative Change |
|--------------------|----------|---------|-----------------|-----------------|
| Subject A Placebo  | 15.7%    | 15.6%   | -0.04%          | -0.3%           |
| Subject B 10 mg QD | 24.6%    | 6.0%    | -18.6%          | -75.6%          |



Subject A, Segment 8

Baseline

Week 12




Subject B, Segment 8



# VK2809 Cohorts Demonstrated High Relative Response Rates


- ▶ Up to 100% of VK2809 patients experienced response, as defined by  $\geq 30\%$  decrease in liver fat at Week 12
- ▶ Combined VK2809 cohorts demonstrate 88% response rate
- ▶ 70% of all patients receiving VK2809 demonstrated liver fat reductions  $\geq 50\%$
- ▶ Reduction in liver fat correlated with improved odds of long-term histology benefit<sup>1</sup>

## Patients with $\geq 30\%$ Relative Reduction in Liver Fat at 12 Weeks




# VK2809 Improved Atherogenic Protein Levels at 12 Weeks

Mean Change in Lipoprotein(a) at Week 12



Mean Change in Apolipoprotein B at Week 12



- ▶ Reductions in Lp(a), ApoB achieved at 12 Weeks
- ▶ Suggests potential cardiovascular benefit

# VK2809-201: Summary of Safety and Adverse Events

|                                                            | Placebo | VK2809<br>5 mg QD | VK2809<br>10 mg QOD | VK2809<br>10 mg QD | All<br>VK2809 |
|------------------------------------------------------------|---------|-------------------|---------------------|--------------------|---------------|
| Randomized (n)                                             | 17      | 10                | 16                  | 16                 | 42            |
| Number (%) of subjects completing study                    | 11 (65) | 9 (90)            | 11 (69)             | 12 (75)            | 32 (76)       |
| Number (%) of subjects with treatment-emergent AEs (TEAEs) | 9 (53)  | 6 (60)            | 10 (63)             | 10 (63)            | 26 (62)       |
| Number of Serious TEAEs                                    | 0       | 0                 | 0                   | 0                  | 0             |
| Number (%) of subjects discontinued due to AE              | 2 (12)  | 1 (10)            | 3 (19)              | 1 (6)              | 5 (12)        |
| Number (%) of subjects with CV-related AEs                 | 3 (18)  | 0                 | 3 (19)              | 0*                 | 3 (7)         |
| Number (%) of subjects with diarrhea or nausea             | 3 (18)  | 1 (10)            | 0                   | 2 (13)             | 3 (7)         |

\*Notes: One subject reported SVT at Week 16 visit, not during 12 week dosing period

- ▶ Encouraging safety and tolerability, no dose-related trends
- ▶ No serious adverse events observed in any arm
- ▶ Excellent GI tolerability

# VK2809-201: Safety Summary

- ▶ Adverse events relatively evenly distributed across placebo, treatment groups
- ▶ Mean ALT, AST levels in VK2809-treated subjects were reduced relative to placebo at Week 12
  - Patients with elevated baseline ALT demonstrated greater improvement relative to placebo at Weeks 12 and 16
- ▶ No other liver function tests were significantly different from placebo at Week 12
- ▶ No clinically meaningful changes were noted among VK2809-treated patients relative to placebo-treated patients for:
  - Thyroid hormones (fT4, tT3, TSH)
  - Cardiovascular safety markers (troponin, CK-MB, NT proBNP)
  - Vital signs (systolic BP, diastolic BP, heart rate, weight)

# VK2809-201: Summary and Conclusions

- ▶ VK2809 produced robust reduction in liver fat on MRI-PDFF in NAFLD patients after 12 weeks of oral dosing
- ▶ 88% of patients receiving VK2809 experienced  $\geq 30\%$  reduction in liver fat content at 12 weeks, including all patients receiving 5 mg daily doses
- ▶ 70% of VK2809 treated patients experienced liver fat reductions  $\geq 50\%$
- ▶ VK2809 produced significant reduction in LDL-C, triglycerides, Apo B, and Lp(a), suggesting potential long-term CV benefit
- ▶ VK2809 was safe in this 12 week Phase 2 study; No SAEs observed
- ▶ VK2809 was well-tolerated in this 12 week Phase 2 study
  - Diarrhea, nausea rates higher in placebo (18% vs. 7% for combined VK2809 cohorts)

## VK2809-201: Next Steps

- ▶ Submit IND to GI division 2H19
- ▶ Initiate Phase 2b study in biopsy-confirmed NASH



# Rare Disease Program

VK0214


X-Linked Adrenoleukodystrophy

- ▶ X-Linked adrenoleukodystrophy (X-ALD)
  - Orphan neurodegenerative disorder
  - X-linked: Carried by females, primarily manifesting in males
  - No cure, no approved therapy
- ▶ Most severe form: Cerebral ALD
  - Rapidly progressive inflammatory demyelination; disruption of BBB
  - Affects ~35% before age 12 (CCALD), ~20% between age 20 – 35 (CALD)
  - Deterioration in speech, cognition; vegetative state within 3-5 years
- ▶ Most common form: Adrenomyeloneuropathy (AMN)
  - Affects spinal cord, motor neurons; no inflammatory component or brain involvement
  - Affects nearly all adult patients; considered “default” manifestation of ALD
  - Progressive motor impairment; wheelchair confinement, leg paralysis common

# TR $\beta$ : X-Linked Adrenoleukodystrophy

Caused by mutation in gene for the ATP-Binding Cassette transporter D1 (ABCD1)

- ▶ Peroxisomal transporter of very long chain fatty acids (VLCFA)



**ABCD1:** Normal function to transport VLCFA into peroxisome for degradation

**X-ALD:** Defective ABCD1 leads to accumulation of VLCFA in tissues


**High VLCFA levels** disrupt cell membranes; inflammatory demyelination in brain tissue; motor neuron deterioration

**TR $\beta$  Agonists:** Stimulate expression of compensatory transporters ABCD2, 3; may mitigate VLCFA elevation

# VK0214: *In Vivo* Proof-of-Concept Data, ABCD1 KO Mouse

- ▶ ABCD1 Knockout model: Mimics biochemical features of human X-ALD
- ▶ VK0214: Durable and progressive reductions in plasma VLCFAs
  - Tissue effects suggest encouraging CNS activity following long-term exposure


## Reductions in Plasma VLCFA-LPC, ABCD1 Knockout Model



# VK0214: Reduces VLCFA Levels in Key Tissues

- ▶ Significant VLCFA reductions observed in multiple tissues
- ▶ Encouraging evidence of CNS activity
- ▶ Reductions in multiple VLCFAs consistent with plasma observations
- ▶ Suggests potential benefit in both cerebral and AMN forms of X-ALD
- ▶ Next steps: IND-filing, planned 2H19

## Change in Tissue VLCFAs: CNS and Peripheral Tissue



|               |       |       |         |      |
|---------------|-------|-------|---------|------|
| % Difference: | -19%  | -15%  | -34%    | -11% |
| p-value:      | <0.05 | <0.01 | <0.0001 | 0.07 |



# Musculoskeletal Program

VK5211: Selective Androgen Receptor Modulator

Hip Fracture

# Musculoskeletal Program: VK5211 for Hip Fracture

- ▶ **Novel Selective Androgen Receptor Modulator (SARM), for hip fracture**
  - VK5211 is a potentially best-in-class small molecule SARM
- ▶ **Promising efficacy signals to date**
  - Significant improvements in lean body mass in animals and humans
- ▶ **Target indication: Rehabilitation post-hip fracture**
  - Population rapidly loses muscle, bone, function
  - >300,000 patients per year, U.S.; expect continued growth
  - Market opportunity potentially exceeds \$1B annually
- ▶ **Potential benefits in other fractures, joint replacements, and muscle wasting disorders**



# VK5211: Potential Therapeutic Benefits of SARMs



- ↑ Muscle Mass
- ↑ Muscle Strength




- ↑ Bone Formation
- ↑ Bone Strength
- ↓ Bone Resorption



- ↑ Cognition
- ↑ Libido
- ↑ Energy

Goal: Retain beneficial properties of natural androgens with fewer side effects relative to anabolic steroids

# VK5211: Phase 2 Trial, Hip Fracture




## ► 12-week Phase 2 trial in 108 patients

- Once-daily oral dosing
- Primary endpoint: Change in lean body mass
- Secondary and exploratory endpoints: Change in appendicular lean mass, total lean body mass, BMD, functional status, ADL, QOL

# Phase 2 Results Show Significant Increases in Lean Body Mass


- ▶ Significant increases in LBM and appendicular lean mass following 12 weeks of daily dosing
- ▶ Consistent dose response observed across primary, secondary efficacy measures
- ▶ Encouraging safety and tolerability, no drug-related SAEs



VK5211 produced robust dose-dependent effects on primary and all secondary measures of lean body mass


# VK5211-201: Change in Body Composition at 12 Weeks

Change in Mean Body Weight, at Week 12



|         |         |         |         |         |
|---------|---------|---------|---------|---------|
| Change  | 0.70 kg | 2.54 kg | 2.95 kg | 3.09 kg |
| p-value | -       | NS      | NS      | NS      |

Change in Mean Fat Mass, at Week 12




|         |      |      |       |       |
|---------|------|------|-------|-------|
|         | 2.9% | 0.7% | -2.5% | -3.3% |
| p-value | -    | 0.24 | 0.11  | 0.01  |

- ▶ Dose-dependent increase in mean body weight
- ▶ Dose-dependent decrease in mean fat mass; significant at high dose

# VK5211 Demonstrates Potency Advantages to Other Mechanisms

- ▶ Significantly greater potency relative to testosterone analogs
- ▶ Improved potency on muscle compared with myostatin-targeting approaches
- ▶ Competitive effect on total lean body mass relative to other clinical-stage SARMs

## Reported Changes in Total Lean Body Mass in Older Adults



|               |        |        |        |       |      |        |
|---------------|--------|--------|--------|-------|------|--------|
| % Difference: | 6.3%   | 8.2%   | 9.9%   | 0.8%  | 1.2% | 2.8%   |
| p-value:      | <0.005 | <0.001 | <0.001 | <0.05 | 0.02 | <0.001 |

VK5211 in older adults provides increases in lean body mass that compare favorably to other developmental mechanisms

# VK5211 Phase 2 Study: Overall Safety and Adverse Events

- ▶ Encouraging safety and tolerability
- ▶ Frequency of reported adverse events demonstrates no dose-relationship
- ▶ No drug-related SAEs
- ▶ Next steps
  - Exploring collaboration and licensing opportunities
  - Plan for next steps with partner

# Financial Summary

## ► Capital structure and summary financials

| Capital Structure <sup>1</sup>        | In '000s |
|---------------------------------------|----------|
| Shares outstanding                    | 72,028   |
| Options, RSUs                         | 3,067    |
| Warrants                              | 6,130    |
| Total shares, options, RSUs, warrants | 81,225   |

| Financials                      | March 31, 2019<br>(\$'000s) |
|---------------------------------|-----------------------------|
| Cash burn 1Q 2019               | \$3,152                     |
| Cash and ST Investments         | \$298,718                   |
| Notes: 1) As of March 31, 2019. |                             |

# Investment Highlights

## ► Focused on best-in-class drugs for metabolic and endocrine diseases

- Two Phase 2 programs supported by positive clinical data

## ► Metabolic Disease Program: VK2809 for NASH

- Novel, selective thyroid receptor- $\beta$  (TR $\beta$ ) agonist
- Phase 2 results demonstrate significant reduction in liver fat content, lipids

## ► Rare Disease Program: VK0214 for X-ALD

- Second selective thyroid receptor- $\beta$  agonist
- In vivo data show improvement in key biomarkers

## ► Musculoskeletal Program: VK5211 for hip fracture recovery

- Non-steroidal selective androgen receptor modulator (SARM)
- Phase 2 results demonstrate significant increases in lean body mass

The background of the slide features a blurred image of laboratory glassware, including several glass bottles and test tubes, overlaid with a blue geometric design consisting of diagonal stripes.

NOVEL THERAPEUTICS  
FOR **METABOLIC &**  
**ENDOCRINE DISORDERS**

Corporate Presentation